


Unit 1: BASIC CHEMISTRY

Chapter 1: Chemical Foundations

1.4: Uncertainty in Measurement

Uncertainty: - all measuring instruments have uncertainty due to how the instrument was manufactured or reading error by the user.



1.5: Scientific Notations and Significant Notations

Scientific Notation: - commonly used to state very big or very small numbers.

$$(1 \text{ to } 9.999\dots) \times 10^n$$

n is an integer

If $n < 0$, then the actual number was smaller than 1

If $n > 0$, then the actual number was greater than 10

Example 1: Convert the following standard notations to scientific notations or vice versa.

- Speed of Light = 3×10^5 km/s = **300,000 km/s** (moved 5 decimal places to the right)
- Mass of an Electron = 9.11×10^{-31} kg = **0.000 000 000 000 000 000 000 000 000 911 kg** (moved 31 decimal places to the left)
- Diameter of a Red Blood Cell = 0.000 007 5 m = **7.5×10^{-6} m** (moved 6 decimal places to the right)
- 2003 US Debt = \$6,804,000,000,000 = **$\6.804×10^{12}** (moved 12 decimal places to the left)

Significant Digits: - digits used in the measurement plus one uncertain value.

Exact Number: - number that indicates no uncertainty. (Numbers in formulas; numbers written in words)

To Count Significant Digits

1. Start counting the first non-zero digit. Do NOT count the leading zero(s).
2. Count all **captive zeros** (between non-zero digits) and **trailing zero** at the end of the measurement.

Example 2: State the number of significant digits for the following measurements.

- | | | | |
|--------------|----------------------|----------------------------|----------------------|
| a. 0.03 g | 1 significant digit | e. 25 000 g | 5 significant digits |
| b. 0.030 g | 2 significant digits | f. 9.300×10^4 m | 4 significant digits |
| c. 0.0304 g | 3 significant digits | g. 4.05×10^{-2} L | 3 significant digits |
| d. 0.03040 g | 4 significant digits | | |

Calculating with significant digit

1. **Adding and Subtracting:** - Line up the significant digits. The answer should be to the least precise measurement used in the calculation.

Example 3: $5.345 \text{ g} + 2.42 \text{ g} + 1.8 \text{ g}$

$$\begin{array}{r} 5.345 \\ 2.42 \\ +1.8 \\ \hline 9.565 \end{array}$$

Least precise decimal place is at the tenth. Final Answer should be to one decimal place.

9.6 g


2. **Multiplying and Dividing:** - answer should be in the least number of significant digits used in calculation.

Example 4: $\frac{3.250 \text{ g}}{1.4 \text{ mL}}$

$$\frac{3.250 \text{ g}}{1.4 \text{ mL}} = 2.321428571 \text{ g/mL}$$

The least number of significant digits used is two.

2.3 g/mL




3. **Multiple Step Calculations:** - follow the multiply and divide rule.
- **Do NOT round off until the very LAST step.**

Example 5: Calculate the final output energy in *Joules* if the equivalent mass of 5.3×10^{-3} kg is turned into energy along with an initial energy input of 4.15×10^{14} J.

(Use $E = mc^2$ where $c = 3.00 \times 10^8$ m/s) **Scientific Notation ($\times 10^n$)**

$$\begin{aligned} E_{\text{output}} &= mc^2 + E_{\text{input}} \\ E_{\text{output}} &= (5.3 \times 10^{-3} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2 + 4.15 \times 10^{14} \text{ J} \\ E_{\text{output}} &= 8.92 \times 10^{14} \text{ J} \end{aligned}$$

$E_{\text{output}} = 8.9 \times 10^{14} \text{ J}$



2nd

EE

Theoretical Result: - the supposed result of an experiment according to pre-lab calculation.

Experimental Result: - the actual measured result of an experiment.

Example 6: Determine the % Error and % Yield of an experiment if the theoretical result was 4.579 g and the experimental result was 4.272 g.

$$\% \text{ Error} = \frac{|\text{Theoretical} - \text{Experimental}|}{\text{Theoretical}} \times 100\% \qquad \% \text{ Yield} = \frac{\text{Experimental}}{\text{Theoretical}} \times 100\%$$

(100% is an exact number)

$$\% \text{ Error} = \frac{|4.579 \text{ g} - 4.272 \text{ g}|}{4.579 \text{ g}} \times 100\%$$

$$\% \text{ Error} = 6.705\%$$

$$\% \text{ Yield} = \frac{4.272 \text{ g}}{4.579 \text{ g}} \times 100\%$$

$$\% \text{ Yield} = 93.30\%$$

1.3: Units of Measurement

SI Units: - International Metric Units (*le Système International*).

Metric Prefixes and Exponential Notations

| Giga | ** | Mega | ** | kilo | hecto | deca | Basic Units | deci | centi | milli | ** | micro | ** | nano |
|--------|----|--------|----|--------|--------|--------|-------------|-----------|-----------|-----------|----|-----------|----|-----------|
| G | | M | | k | h | da | metre (m) | d | c | m | | μ | | n |
| 10^9 | | 10^6 | | 10^3 | 10^2 | 10^1 | Litre (L) | 10^{-1} | 10^{-2} | 10^{-3} | | 10^{-6} | | 10^{-9} |
| | | | | | | | gram (g) | | | | | | | |
| | | | | | | | Kelvin (K) | | | | | | | |
| | | | | | | | Pascal (Pa) | | | | | | | |
| | | | | | | | Newton (N) | | | | | | | |
| | | | | | | | Mole (mol) | | | | | | | |

When converting units, the same number of significant digits must be preserved.

$$\begin{array}{ccc} & \times 1000 & \\ & \curvearrowright & \\ 1 \text{ mL} & = & 1 \text{ cm}^3 \\ & \curvearrowleft & \\ 1 \text{ L} & = & 1000 \text{ cm}^3 \\ & \times 1000 & \end{array}$$

Note: $1 \text{ m}^3 \neq 1 \text{ L}$

$$1 \text{ m}^3 = 1 \text{ m} \times 1 \text{ m} \times 1 \text{ m}$$

$$1 \text{ m}^3 = 100 \text{ cm} \times 100 \text{ cm} \times 100 \text{ cm}$$

$$1 \text{ m}^3 = 1,000,000 \text{ cm}^3$$

$$1 \text{ m}^3 = 1000 \text{ L}$$

Example 1: Complete the following unit conversions

- a. 345 mL = **0.345** L (left 3 places) d. 26 cm³ = **0.026** L (26 cm³ = 26 mL) (left 3 places)
- b. 42 g = **0.042** kg (left 3 places) e. 1854 cm = **0.01854** km (left 5 places)
- c. 54300 m = **54.300** km (left 3 places) f. 0.035 kg = 35000 mg = **3.5 × 10⁴** mg (right 6 places)

too many significant; original measurement only has two digits.

two significant digits

Assignment

1.4 pg. 32 #19, 20

1.5 pg. 33 #23, 25 to 32

1.6: Dimensional Analysis

Dimensional Analysis: - commonly known as unit factor method.

- using units to analyse unit conversion or whether the right kind of procedure is used for calculations.
- unit factors have 1 bigger unit along with equivalent smaller unit.
- should keep the original number of significant digits.

Example 1: Convert 65.0 miles/h to km/h. (1 mile = 1.609344 km)

$$\frac{65.0 \text{ miles}}{1 \text{ hour}} \times \frac{1.609344 \text{ km}}{1 \text{ mile}} = 104.60736 \text{ km/h} \quad (\text{round to 3 significant digits})$$

105 km/h

Example 2: Convert 50 km/h to m/s.

$$\frac{50 \text{ km}}{1 \text{ hour}} \times \frac{1000 \text{ m}}{1 \text{ km}} \times \frac{1 \text{ hour}}{3600 \text{ s}} = 13.8888889 \text{ m/s} \quad (\text{round to 2 significant digits})$$

14 m/s

Example 3: Convert 55 miles/gal to km/L. (1 gal = 3.785412 L)

$$\frac{55 \text{ miles}}{1 \text{ gal}} \times \frac{1.609344 \text{ km}}{1 \text{ mile}} \times \frac{1 \text{ gal}}{3.785412 \text{ L}} = 23.38290257 \text{ km/L} \quad (\text{round to 2 significant digits})$$

23 km/L

1.7: Temperature

Temperature: - the average kinetic energy of a substance.

$T_F = \frac{9}{5}T_C + 32$

$T_F - 32 = \frac{9}{5}T_C$

$\frac{5}{9}(T_F - 32) = T_C$

Convert from degree Celsius to Fahrenheit

32 is an exact number

Convert from Fahrenheit to degree Celsius

Some important temperatures:

0°C = Water Freezes
 100°C = Water Boils

37°C = Normal Body Temperature
 20°C = Ambient Room Temperature

Kelvin: - temperature scale where **0 K (absolute zero) = -273.15°C** (freezing point of hydrogen – no heat, particles stop moving)

$$T_K = T_C + 273.15$$

Example 1: With wind chill, Calgary can get down to -37°C . Convert the temperature to Fahrenheit and Kelvin.

$$T_F = \frac{9}{5}T_C + 32$$

$$T_K = T_C + 273.15$$

$$T_F = \frac{9}{5}(-37) + 32$$

$$T_K = -37 + 273.15$$

$$T_F = -34.6^\circ\text{F}$$

$$T_K = 236.15\text{ K}$$

$$T_F = -35^\circ\text{F}$$

$$T_K = 236\text{ K}$$

Assignment

1.6 pg. 33–34 #33, 35 (refer to Table 1.4 on pg. 18), 41, 42, 43, 45

1.7 pg. 34–35 #47, 49, 51

1.8: Density**Mass:** - the amount of stuff in an object.**Weight:** - the amount of gravitational force that is pulling on an object.

Example: An object that has a mass of 50 kg on Earth will have a mass of 50 kg on the moon. However, the same object, which has a weight of 490.5 N on Earth, will only weight 81.75 N on the moon. This is because the gravitation pull on the moon is 1/6 of that on Earth.

Density: - the amount of mass per unit of volume

$$\text{Density} = \frac{\text{Mass (g or kg)}}{\text{Volume (cm}^3, \text{ mL, L, m}^3)} \quad D = \frac{m}{V}$$

Example 1: Lead has a density of 11.34 g/cm³. If a lead sphere has a radius of 5.00 cm, what is its mass?

$$D = 11.34 \text{ g/cm}^3 \quad r = 5.00 \text{ cm} \quad m = ?$$

Manipulate the formula to solve for m :

$$D = \frac{m}{V}$$

$$DV = m$$


We need Volume of the sphere.

$$V_{\text{sphere}} = \frac{4}{3}\pi r^3$$

$$V_{\text{sphere}} = \frac{4}{3}\pi(5.00 \text{ cm})^3$$

$$V_{\text{sphere}} = 523.5987\dots \text{ cm}^3$$

(Do NOT round off. We are not done yet.)




Substitute D and V to solve for m

$$m = DV$$

$$m = (11.34 \text{ g/cm}^3)(523.5987\dots \text{ cm}^3)$$

$$m = 5.94 \times 10^3 \text{ g or } 5.94 \text{ kg}$$

To recall all digits of the previous answer



2nd**ANS****(-)**


1.9: Classification of Matter

All substance in the universe is made up of **matter**. However, matter can be classified into **mixtures** and **pure substances**.

There are two kinds of mixtures. **Heterogeneous (*hetero* means different) mixtures** are mechanical mixtures which we can see its different components with the naked eye. An example of a heterogeneous mixture is a bag of assorted nuts. We can clearly see the different kind of nuts (walnuts, peanuts, chestnut, hazelnut ... etc.) in this bag. A **homogeneous (*homo* means the same) mixture** is called a solution. Unlike heterogeneous mixture, a solution is a mixture that consists of different components, which cannot be seen from a naked eye. An example of a solution is a salt solution. After we completely dissolved the salt in water, we cannot see the salt particles in the water.

Unlike mixtures, **pure substance** is a substance with a constant composition that cannot be separated by physical means. Pure Substances can be classified into **elements** and **compounds**. Element is a pure substance that has one kind of **atom**. The Periodic Table of Elements lists all the different elements that are either found in nature or prepared in the laboratory synthetically. An atom is defined as the smallest particle of matter. An example of an element is hydrogen. It contains only hydrogen atom. A compound is defined as a pure substance that is composed of two or more different elements. The smallest unit of a compound is called a **molecule (a particle that is made up of two or more different atoms)**. An example of a compound is water. The smallest unit of water is the H₂O molecule. Each water molecule (H₂O) contains 2 hydrogen atoms and an oxygen atom.

The classification of matter is explained in a flow chart below.




Matter: - anything that has a mass and occupies space.

Mixtures: - are matters that are made up of more than one kind of substances and the components can be separated by **physical change (change of state, stirring, filtering... etc)**.

Heterogeneous (Mechanical) Mixture: - mixture that is composed of two or more substances where the components can be seen by the naked eye.

Filtration: - using a filter and a funnel, a mechanical mixture consists of a liquid can be separated.



Homogeneous Mixture (Solution): - mixture that is composed of two or more substances where the components the same throughout (cannot separate the components by the naked eye).

Solute: - the substance being dissolve.

Solvent: - the substance doing the dissolving

Example: Salt Water (Solute = Salt; Solvent = Water) 9% Alcohol (Solute = Alcohol; Solvent = Water)

Evaporation: - an aqueous solution that consists of a solid solute can be recovered by evaporation of the solvent. The solvent may be recovered as well if a condensation device is used.

Distillation: - an aqueous solution that consists of a liquid solute can be separated by evaporation of the substance with a lower boiling point followed by condensation.

Pure Substance: - a substance with a constant composition.

- in a case where the pure substance is composed of more than one kind of matter, they can only be separated by **chemical change (burning, oxidation, electrolysis ... etc)**.

Element: - a pure substance that is made up of one kind of atom.

Compound: - a pure substance that is made up of more than one kind of element.

Atom: - the smallest particle of matter.

Molecule: - the smallest unit of a compound (a particle that is made up of more than one kind of atom).

Examples:

1. Classify the following as Heterogeneous or Homogeneous Mixture:

- a) a bag of gravel b) cement c) saturated salt water d) a methanol and water mixture
 e) oil and water f) the atmosphere g) Jell-O h) diet carbonated soft drink

2. Classify the following as Mixture or Pure Substance:

- a) lake water b) tap water c) distilled water d) iron e) steel (iron and carbon)
 f) chromium g) beer h) sugar i) gasoline

3. Classify the following as Element or Compound: (use the Periodic Table of Elements)

- a) hydrogen e) carbon dioxide i) salt m) propane
 b) water f) chlorine j) nickel n) baking soda
 c) ammonia g) ethanol k) gold o) uranium
 d) oxygen h) charcoal (carbon) l) neon p) mercury

Answers:

1. Heterogeneous Mixtures: a), e)

Homogeneous Mixtures: b), c), d), f), g), h)

All the components of the heterogeneous mixtures can be seen by the naked eye. However, the components of the homogeneous mixtures cannot be distinguished by the naked eye.

2. Mixtures: a) lake water: contains water, soil particles, micro-organisms ...etc.

b) tap water: contains fluoride and chloride additives.

e) steel: a mixture of iron and carbon.

g) beer: contains alcohol, water and other ingredients.

i) gasoline: contains

Pure Substances: c) distilled water: contains water (H₂O) only.

d) iron: an element with a symbol Fe.

f) chromium: an element with a symbol Cr.

h) sugar: a compound commonly known as sucrose (C₁₂H₂₂O₁₁).

3. Elements: a) hydrogen (H)

d) oxygen (O)

f) chlorine (Cl)

h) carbon (C)

j) nickel (Ni)

k) gold (Au)

l) neon (Ne)

o) uranium (U)

p) mercury (Hg)

Compounds: b) water (H₂O)

c) ammonia (NH₃)

e) carbon dioxide (CO₂)

g) ethanol (C₂H₅OH)

i) salt (NaCl)

m) propane (C₃H₈)

n) baking soda (NaHCO₃)

If the name of the substance appears on the Periodic Table of Elements, then it is an element.

Assignment


1.8 pg. 35 #55, 57, 59, 61, 63

1.9 pg. 35–36 #65, 67, 69

Chapter 2: Atoms, Molecules and Ions**2.2: Fundamental Chemical Laws**

1. **Law of Conservation of Mass:** - mass is neither created nor destroyed in a chemical reaction
(Lavoisier)

Example 1: A 3.50 g of iron nail is allowed to rust. The rusted nail has a mass of 5.1 g. What is the amount of oxygen reacted with the iron nail?



2. **Law of Definite Proportion:** - the same compound always contains exactly the same proportion of elements by mass.
(Proust)

Example 2: Water contains about 8 parts oxygen to 1 part hydrogen by mass.

3. **Law of Multiple Proportion:** - when two elements form a series of compounds, the ratios of the masses of the second element that combine with the first element can always be reduced to small whole numbers.
(Dalton)

Example 3: State the ratios of hydrogen between the following hydrocarbon compounds.

| Hydrocarbons | Mass of Hydrogen per 1 g of Carbon |
|--------------|------------------------------------|
| Compound A | 2.973 g |
| Compound B | 3.963 g |
| Compound C | 4.459 g |

| | |
|---|---|
| $\frac{A}{B} = \frac{2.973 \text{ g}}{3.963 \text{ g}} = 0.7501892506 \approx 0.75$ | $\frac{A}{B} = \frac{3}{4}$ $\frac{B}{C} = \frac{8}{9}$ $\frac{C}{A} = \frac{3}{2}$ |
| $\frac{B}{C} = \frac{3.963 \text{ g}}{4.459 \text{ g}} = 0.8887642969 \approx 0.888\dots$ | |
| $\frac{C}{A} = \frac{4.459 \text{ g}}{2.973 \text{ g}} = 1.49983182 \approx 1.5$ | |

2.3: Dalton Atomic Theory

The practice of using symbols to represent elements can be traced back to the ancient Greek alchemists. Their purpose was to find a chemical recipe to make gold from other less valuable metals. (We now know that it is only possible now if we can change the number of protons in the nucleus).

In 1808, a British scientist by the name of John Dalton published his theory of atoms that would have profound effects on the philosophy on chemistry and physics. (The word atom comes from the Greek word *atomos*, which means indivisible. A Greek philosopher Democritus in 5th-century BC first suggested this concept). The **Dalton's Atomic Theory** can be summarized as:

- 1. All elements are made up of tiny particles called atoms.**
- 2. The atoms of a particular element are identical. Different elements have different kind of atoms.**
- 3. Atoms cannot be created or destroyed.**
- 4. Chemical compounds are formed when different kinds of atoms combine together. A particular compound always has the same relative numbers and types of atoms.**
- 5. Chemical reactions deal with the rearrangement of the atom, which changes the way they are combined together. There is no change to the atoms themselves in a chemical reaction.**





Today, all elements and their symbols are listed in an orderly fashion with the help of the Periodic Table of Elements. The use of standardised symbols allows scientists from all over the world to share their knowledge despite the differences in language. Most elements are **monoatomic**. That means their atoms can exist individually ("*mono*" means one). Others are **diatomic**, atoms that exist in pairs ("*di*" means two). Some are **polyatomic**, atoms that exist in numbers more than one ("*poly*" means many). The table on the next page shows all the diatomic and polyatomic elements.


| Diatomic Elements (all the ~gens, including Halogens - second last column of the Periodic Table) | Polyatomic Elements | Monoatomic Elements |
|---|---|---|
| Hydrogen (H ₂), Oxygen (O ₂), Nitrogen (N ₂), Fluorine (F ₂), Chlorine (Cl ₂), Bromine (Br ₂), Iodine (I ₂), Astatine (At ₂) | Phosphorus (P ₄) Sulphur (S ₈) | All other Elements. Examples: Helium (He), Iron (Fe), Calcium (Ca), Silver (Ag), Mercury (Hg) |

Note: **Students should memorise all the diatomic and polyatomic elements. They are the only exceptions. All other elements are monoatomic.** A lot of the symbols are recognisable from the name of the elements (Zinc: Zn; Carbon: C; Aluminium: Al). Some of them look somewhat different. This is because the symbols came from the elements' Latin names (Silver: Ag for "Argentum"; Gold: Au for "Aurum"). To save time, students should also familiarise themselves with the whereabouts of the elements on the Table.

2.4: Early Experiments to Characterise the Atom

Since the time of Dalton's Atomic Theory, scientists had improved upon his model to better explain the structure of an atom. The following is a summary of the different atomic models.

| | |
|---|---|
|  | <p><u>Dalton's Atomic Model:</u></p> <p>In 1808, John Dalton proposed that all matter is made up of tiny particles called atoms. Atoms cannot be divided, created or destroyed. Different elements have different kinds of atoms. The difference is mainly due to the different atomic masses.</p> |
|  <p>Positive Charged Spherical Cloud</p> | <p><u>Plum Pudding Model:</u></p> <p>In 1903, J.J. Thomson and Michael Faraday discovered electrons within an atom using a device called the cathode ray tube. Electrons are negatively charged subatomic particles with a charge of -1. The electrons were viewed as embedded in a positively charged spherical cloud. This is similar to the raisins distributed in a plum pudding. Robert Millikan used his oil drop experiment (by balancing the weight of an oil drop with electric force) to determine the elemental charge of the electron as 1.6×10^{-19} C and has a mass of 9.11×10^{-31} kg.</p> |
|  <p>Electrons</p> <p>Nucleus (Protons and Neutrons)</p> | <p><u>Nuclear Model:</u></p> <p>In 1912, Ernest Rutherford proposed the Nuclear Model for atoms after his famous gold foil experiment. Earlier to this time, E. Goldstein discovered the positively charged ($+1$) subatomic particles called protons. Rutherford proposed that the protons are packed tightly together at the centre of the atom called the nucleus. In 1932, James Chadwick discovered neutrons (no charged). Together, they suggested that the nucleus was made up of both protons and neutrons (the bulk of the atomic mass) since electrons are very light compared to the masses of the protons and neutrons.</p> <p>On the other hand, negatively charged electrons move around the nucleus because of their attraction with the positively charged nucleus (contains protons). Since the nucleus is very small, the circling electrons make up almost all of the volume of the atom. If the atom has a size of a football field, the nucleus is about the size of a small nail at the centre of the field.</p> |
|  <p>Orbits – Energy Levels</p> | <p><u>The Bohr Model:</u></p> <p>In 1913, Neil Bohr refined the Nuclear Model by suggesting that electrons move around the nucleus in specified orbits. These orbits are called energy levels. Electrons cannot exist between the orbits. The further the orbit is from the nucleus, the higher its energy level for the electrons in that orbit. This is very similar to the planetary model of our Solar system.</p> |



“Electron Clouds”
Orbitals

The Electron Cloud (Quantum Mechanics) Model:

This modern atomic model is very similar to the Bohr model. We still use the energy levels, however, the idea of orbits is modified into **orbitals**. **An orbital is a region of space where the electrons are most probably in.** Calculations of these orbital shapes involve advanced mathematics. Scientists use this model with the Molecular Orbital Theory to predict complex reactions and possible new chemical compounds.

2.5: Modern View of Atomic Structure

| Subatomic Particles | Charge | Relative Mass | Actual Mass | Location |
|-------------------------------------|--------|---------------|---------------------------|--------------------------------------|
| Electrons (e^-) | -1 | 1 | 9.11×10^{-31} kg | Region around the center of the atom |
| Protons (p^+) | +1 | 1836.12 | 1.67×10^{-27} kg | Centre of the atom called Nucleus |
| Neutrons (n) | 0 | 1836.65 | 1.67×10^{-27} kg | Inside the Nucleus with the protons |

Atomic Number and Atomic Mass:

We have looked at different atomic models. In high school chemistry, we deal mainly with the Bohr model. Recall from the Dalton's Atomic Theory, one of its points is that different elements have different atoms. The main difference between them is the **atomic mass**. This is the mass characteristic of a given element. **The atomic mass of an element is relative to the mass of the carbon atom (6 protons and 6 neutrons with an atomic mass of 12).** It is usually located at the right, top corner or directly below each element on the Table of Elements. For now, we will assume that atomic mass has a unit of amu (Atomic Mass Unit). We will revisit the atomic mass in Section 1.6.

Because different elements have different atomic mass, the number of subatomic particles within an atom is also different for these elements. The **atomic number, a number assigned to each element based on its atomic mass**, is located at the top left corner of each element on the Table of Elements. **The atomic number is equated to the number of protons and electrons of that atom.** The **number of neutrons can be found by subtracting the atomic mass (rounded off whole number) with the atomic number.**

Atomic Number = Number of Protons and Electrons of an Atom

Number of Neutrons = Atomic Mass – Atomic Number

Example 1: State the Atomic Number, Atomic Mass, number of protons, neutrons, and electrons of the following elements.

| | |
|-----------|-------|
| 17 | 35.45 |
| Cl | |
| Chlorine | |

Atomic Number = 17
 (17 p^+ and 17 e^-)
Atomic Mass = 35.45
 # of Neutron = $35.45 - 17 = 18$ n

| | |
|----------|------|
| 1 | 1.01 |
| H | |
| Hydrogen | |

Atomic Number = 1
 (1 p^+ and 1 e^-)
Atomic Mass = 1.01
 # of Neutron = $1.01 - 1 = 0$ n

Note: Because any given atom has the same number of protons and electrons (same atomic number), **all Atoms have a Net Charge of 0.**


Electron Shells, Energy Levels and Valence Electrons:

Recall from the Bohr Model studied earlier. It states that **electrons travel around the nucleus in specified orbits (electrons are quantum)**. These orbits are called **energy levels**. They can also be called **electron shells**. These orbits are very similar to the planets orbiting our sun. The only difference is that each orbit can accommodate more than one electron at a time. The following table shows the maximum number of electrons each successive “orbit” or energy level allows.

| Energy Level | Maximum Number of Electrons Allowed |
|-----------------|-------------------------------------|
| 1 st | 2 |
| 2 nd | 8 |
| 3 rd | 8 |
| 4 th | 18 |
| 5 th | 18 |
| 6 th | 32 |
| 7 th | 32 |

To put electrons in the shells, we have to fill the first energy level until it is full before we can start filling the next energy level. If the second energy level is filled, then we can put electrons in the third energy level and so on. This process is repeated until all the electrons are used up. The following diagrams illustrate the point above.

Valence Electrons: - the electrons in the **outermost** shell.




Element: Hydrogen (H)

Atomic Number: 1 Atomic Mass: 1.01 Nucleus: 1 p⁺ and 0 n

1st Energy Level: 1 e⁻ (1 valence e⁻)
Total: 1 e⁻

Location on the Period Table of Elements: First Row; Column IA




Element: Helium (He)

Atomic Number: 2 Atomic Mass: 4.00 Nucleus: 2 p⁺ and 2 n

1st Energy Level: 2 e⁻ (2 valence e⁻ - Filled)
Total: 2 e⁻

Location on the Period Table of Elements: First Row; Column VIIIA




Element: Lithium (Li)

Atomic Number: 3 Atomic Mass: 6.94 Nucleus: 3 p⁺ and 4 n

2nd Energy Level: 1 e⁻ (1 valence e⁻)
1st Energy Level: 2 e⁻
Total: 3 e⁻

Location on the Period Table of Elements: Second Row; Column IA



Element: Beryllium (Be)


Atomic Number: 4 Atomic Mass: 9.01 Nucleus: 4 p^+ and 5 n

2nd Energy Level: 2 e^- (2 valence e^-)

1st Energy Level: 2 e^-

Total: 4 e^-

Location on the Period Table of Elements: Second Row; Column IIA



Element: Boron (B)


Atomic Number: 5 Atomic Mass: 10.81 Nucleus: 5 p^+ and 6 n

2nd Energy Level: 3 e^- (3 valence e^-)

1st Energy Level: 2 e^-

Total: 5 e^-

Location on the Period Table of Elements: Second Row; Column IIIA



Element: Carbon (C)


Atomic Number: 6 Atomic Mass: 12.01 Nucleus: 6 p^+ and 6 n

2nd Energy Level: 4 e^- (4 valence e^-)

1st Energy Level: 2 e^-

Total: 6 e^-

Location on the Period Table of Elements: Second Row; Column IVA



Element: Nitrogen (N)


Atomic Number: 7 Atomic Mass: 14.01 Nucleus: 7 p^+ and 7 n

2nd Energy Level: 5 e^- (5 valence e^-)

1st Energy Level: 2 e^-

Total: 7 e^-

Location on the Period Table of Elements: Second Row; Column VA



Element: Oxygen (O)


Atomic Number: 8 Atomic Mass: 16.00 Nucleus: 8 p^+ and 8 n

2nd Energy Level: 6 e^- (6 valence e^-)

1st Energy Level: 2 e^-

Total: 8 e^-

Location on the Period Table of Elements: Second Row; Column VIA




Element: Fluorine (F)

Atomic Number: 9 Atomic Mass: 19.00 Nucleus: 9 p⁺ and 10 n

2nd Energy Level: 7 e⁻ (7 valence e⁻)

1st Energy Level: 2 e⁻
Total: 9 e⁻

Location on the Period Table of Elements: Second Row; Column VIIA



Element: Neon (Ne)

Atomic Number: 10 Atomic Mass: 20.18 Nucleus: 10 p⁺ and 10 n

2nd Energy Level: 8 e⁻ (8 valence e⁻ - Filled)

1st Energy Level: 2 e⁻
Total: 10 e⁻

Location on the Period Table of Elements: Second Row; Column VIIIA

One way to remember the maximum number of electrons for each energy level is to look at the Periodic Table of Elements. There are 2 elements in the first row, hence 2 electrons are allowed in the first energy level. There are 8 elements each in the second and third rows, hence 8 electrons are allowed in each of the second and third energy level. This pattern repeats itself for higher energy levels.

Isotopes:

Isotopes are atoms of an element with the same atomic number but a different mass because of a different number of neutrons. For a given mass of substance, there exist a certain percentage of isotopes. Some isotopes are stable. Others are unstable and they go through a decomposition process called **radioactive decay**.




A common example is the isotope $^{14}_6\text{C}$ (Carbon-14: Carbon with an atomic mass of 14 amu, which has 8 n, 6 p⁺ and 6 e⁻). Naturally occur carbon contains 98.9 % of Carbon-12, 0.55% of Carbon-13 and 0.55% of Carbon-14. Chemists, physicists, archaeologists, geologists, and criminologists commonly use the carbon isotope. Because Carbon-14 is unstable and goes through radioactive decay at a definite rate, we can measure the amount of isotopes left in a substance to deduce its age. **Carbon-14 dating is a technique to date archaeological and geological findings by measuring the amount of Carbon-14 left in the artefacts.** Carbon-13 is used by chemists to assist in identifications of various chemical compounds.

Isotopes of other elements also have their uses in society. A tiny proportion of all water molecules (H₂O) composes of a hydrogen isotope called deuterium (^2_1H). Deuterium can be utilised as fuel in nuclear

fusion reactors of the future. Other isotopes of various elements are used as **radiotracers**. These are **radioactive isotopes that can be introduced into organisms in food or drugs, and their pathways can be traced by monitoring their radioactivity**. These radiotracers have found their way into medical research. The list below shows some radiotracers and their medical applications.

| Radiotracers | Area of the body examined |
|---|--------------------------------|
| $^{131}_{53}\text{I}$ | Thyroid |
| $^{59}_{26}\text{Fe}$ and $^{51}_{24}\text{Cr}$ | Red Blood Cells |
| $^{99}_{42}\text{Mo}$ | Metabolism |
| $^{32}_{15}\text{P}$ | Eyes, Liver, Tumours |
| $^{87}_{38}\text{Sr}$ | Bones |
| $^{99}_{43}\text{Tc}$ | Heart, Bones, Liver, and Lungs |
| $^{133}_{54}\text{Xe}$ | Lungs |
| $^{24}_{11}\text{Na}$ | Circulatory System |


Using the same rule, the energy level diagram of an isotope can be drawn as well. Recall that the superscript is the atomic mass of the isotope and the subscript is the atomic number. Since the atomic mass of the isotope is different than the atomic mass of the original element, the number of neutrons of the isotope is different than the number of neutrons of the element (Number of neutron = Atomic Mass – Atomic Number).



Isotope: Tritium (^3_1H)

Atomic Number: 1 Atomic Mass: 3.02 Nucleus: 1 p^+ and 2 n

1st Energy Level: 1 e^- (1 valence e^-)
Total: 1 e^-



Element: Carbon-14 ($^{14}_6\text{C}$)

Atomic Number: 6 Atomic Mass: 14.01 Nucleus: 6 p^+ and 8 n

2nd Energy Level: 4 e^- (4 valence e^-)
1st Energy Level: 2 e^-
Total: 6 e^-

Assignment

2.2 pg. 75–76 #25, 27, 29

2.4 pg. 75–76 #19, 21, 23, 33



2.5 pg. 76 #35 to 46

2.6: Molecules and Ions**Molecules:** - basic unit of a compound.

- contain at least two atoms of the same or different kind of elements.

Ions: - when atoms lose or gain electrons, they attain a positive or negative charge.


- Cations:** - **positive** charged ions (atoms that lose electrons).
- naming cation (element name follow by "ion")

Example 1: Draw the energy level diagrams for the following cations.a. Sodium ion = Na^+ (11 p^+ and 10 e^-)**Sodium Ion (Na^+)****Atomic Number: 11** **Atomic Mass: 22.99** **Nucleus: 11 p^+ and 12 n****2nd Energy Level: 8 e^- (8 valence e^- - Filled)****1st Energy Level: 2 e^-** **Total: 10 e^-** **Net Charge = 1+****Location on the Period Table of Elements: Third Row; Column IA**b. Calcium ion = Ca^{2+} (20 p^+ and 18 e^-)**Calcium Ion (Ca^{2+})****Atomic Number: 20** **Atomic Mass: 40.08** **Nucleus: 20 p^+ and 20 n****3rd Energy Level: 8 e^- (8 valence e^- - Filled)****2nd Energy Level: 8 e^-** **1st Energy Level: 2 e^-** **Total: 18 e^-** **Net Charge = 2+****Location on the Period Table of Elements: Fourth Row: Column IIA**

2. **Anions**: - **negative** charged ions (atoms that gain electrons).
 - naming anion (first part of element name follow by suffix *-ide*)

Example 2: Draw the energy level diagrams for the following anions.

a. Chloride = Cl^- (17 p^+ and 18 e^-)



Chloride (Cl^-)

Atomic Number: 17 **Atomic Mass: 35.45** **Nucleus: 17 p^+ and 18 n**

3rd Energy Level: 8 e^- (8 valence e^- - Filled)

2nd Energy Level: 8 e^-


1st Energy Level: 2 e^-

Total: 18 e^-

Net Charge = 1-

Location on the Period Table of Elements: Third Row; Column VIIA

b. Oxide = O^{2-} (8 p^+ and 10 e^-)



Oxide (O^{2-})

Atomic Number: 8 **Atomic Mass: 16.00** **Nucleus: 8 p^+ and 8 n**

2nd Energy Level: 8 e^- (8 valence e^- - Filled)

1st Energy Level: 2 e^-

Total: 10 e^-

Net Charge = 2-


Location on the Period Table of Elements: Second Row: Column VIA

Octet Rule: - the tendency for electrons to fill the second and third energy levels (8 valence e^-) to achieve stability.

2.7: An Introduction to the Periodic Table

Metals and Non-Metals

The 2 main categories of the Periodic Table of Elements are the **metals** and **non-metals**. They are divided by the “staircase” on the table. This “staircase” can be found at the element Boron extending down to the element Astatine. **Metals are the elements at the left side of the “staircase”, and non-metals are the elements at the right side of the “staircase”.**



Note: Hydrogen is placed at the metal side but is considered a non-metal.

Physical Properties: - are the properties or characteristics of a substance that can be change without involving the chemical change in its composition.

Physical Properties of Metals (with the exception of hydrogen):

1. Metals are mostly solids at room temperature (with the exception of mercury).
2. Metals are malleable (they can be hammered into thin sheets).
3. Metals are ductile (they can be pulled into wires).
4. Metals are good conducts of heat and electricity.
5. Metals are lustrous (shiny).

Physical Properties of Non-Metals:


1. Non-metals are mostly gases or solids at room temperature (with the exception of bromide).
2. Non-metals usually do not have the physical properties of metals.

Some **elements near the “staircase” exhibit both the properties of metals and non-metals.** These elements are referred to as **metalloids**. An example is silicon. It is often used as a semiconductor material (an electrical conductor that can conduct and an insulate electricity). Other metalloids are boron, germanium, arsenic, antimony, tellurium, polonium, and astatine.

Periods and Groups: Chemical Properties of Elements

Chemical Properties: - the properties of a substance that involves a change in the organisation of atoms (mainly the sharing or transfer of electrons).

The shape of the Periodic Table of Elements is a structural way to organise elements. **The vertical columns of the Table are called groups or families.** As we have seen before, the column number is the same as the number of valence electrons of the elements. Since chemical properties depend greatly on the number of valence electrons, all elements within the same group or family must have similar chemical properties. We have already seen one such family, the noble gases. All elements of this group are non-reactive and very stable (recall the valence electron shell of these elements is full). The names of other families and their general chemical properties are listed below.




| Groups or Families | Chemical Properties |
|-----------------------------|---|
| Alkali Metals (IA) | very reactive; forms ions with +1 charge when react with non-metals |
| Alkaline Earth Metals (IIA) | less reactive than alkali metals; forms ions with +2 charge when react with non-metals |
| Halogens (VIIA) | very reactive; form ions with -1 charge when react with metals; all form diatomic molecules |
| Noble Gases (VIIIA) | very stable; do not form ions; monoatomic gas at room temperature |

Periods: - “rows” of elements that are identify by their highest energy level.
 - the pattern of chemical properties “repeats” for every row.

Transition Metals (1B to 10B): - groups and periods of metals that can have **varying charges**.
 - use **Roman Numerals** as part of their ionic names.


Example: Fe^{3+} and Fe^{2+} ions



Chemical Properties of Metals and Non-Metals:

1. **Metals lose electrons to become positive ions – cations.**
2. **Non-Metals gain electrons to become negative ions – anions.**
3. Hydrogen usually loses an electron to become a H^+ ion. However, it can sometimes gain an electron to become H^- (Hydride).
4. **The last column of the Table of Elements does not usually form ions.** These elements are called the **Noble Gases** (Helium, Neon, Argon, Krypton, Xenon, and Radon).

The number of electrons an atom loses or gains depends on which column (vertical) the element is at the Table.



The reason that noble gases (column VIIIA) do not form ions is because their outermost shells are filled with the maximum number of electrons allowed. That is why we call this group of elements “noble gases”. They do not form ions because they are stable. Hence we use the word “noble” to describe them. All the other elements form ions because they want to achieve stability like the noble gases. If you observe carefully, oxide has the same number of electrons as the nearest noble gas, neon. On the other hand, calcium ion has the same number of electrons as the nearest noble gas, argon. In terms of stability, which is another word for lower energy state, these ions are more stable than their respective atoms.


Since the number of valence electrons of an atom is the same as its column number, all the elements of column IA have 1 valence electron. As we see with lithium, all they have to do is to lose that valence electron to achieve a noble gas “like” state. For elements in column IIA, they all have 2 valence electrons. Hence, they lose 2 electrons to acquire stability and become ions with a net charge of +2. The following table summarises these points.

| Column | Number of Valence Electrons | Methods to achieve a Stable State | Net Charge of Ions |
|--------|-----------------------------|--|--------------------|
| IA | 1 | lose 1 electron or gain 7 electrons | +1 |
| IIA | 2 | lose 2 electrons or gain 6 electrons | +2 |
| IIIA | 3 | lose 3 electrons or gain 5 electrons | +3 |
| IVA | 4 | lose 4 electrons or gain 4 electrons | +4 |
| VA | 5 | lose 5 electrons or gain 3 electrons | -3 |
| VIA | 6 | lose 6 electrons or gain 2 electrons | -2 |
| VIIA | 7 | lose 7 electrons or gain 1 electron | -1 |
| VIIIA | 8 | already has the maximum number of electrons allow in the outermost electron shell. | 0 |

Ionic Compound: - when a **metal** combines with a **non-metal**.

- forms **ionic bonds** (electrons are “stolen” or “transfer” from one atom to another).
- dissociates into **electrolytes** (forms ions when dissolve in water)


Example: LiF



Molecular Compound: - when a **non-metal** combines with a **non-metal**.


- forms **covalent bonds** (electrons are “share” between atoms).
- forms **non-electrolytes** (do not dissociate into ions when dissolve in water)

Example: CH₄



Binary Elements: - non-metals that come in pairs of atoms as molecules to achieve stability.

- include all the (*-gens*), hydrogen, nitrogen, oxygen, and all halogens.



Polyatomic Elements: - non-metals that comes in groups of 4 or 8 to achieve stability.

- **P₄ (Phosphorus)** and **S₈ (Sulphur)**

Assignment

2.6 pg. 76 #47, 49

2.7 pg. 76-77 #51, 53 to 55, 57, 58

2.8A: Naming Simple Ionic Compounds**Nomenclature:** - a naming system**IUPAC:** - International Union of Pure and Applied Chemistry.


- an organisation that oversees the standard nomenclature of all chemicals.

Nomenclature of Ionic Compounds


1. Balance the **Cation** and **Anion** Charges.
2. Use **brackets** for **multiple Complex Ions (Polyatomic Ions)**.
3. When naming, use **-ide** for the **non-metal anions**.
4. **Metals** that can have **two or more different charges** must use **Roman Numerals** in the names.

Example 1: Write the chemical formulas for the following.


a. sodium chloride

Need 1 Na^+ & 1 Cl^- to balance charges


b. calcium fluoride

Need 1 Ca^{2+} & 2 F^- to balance charges


c. ammonium sulfate

2 (NH_4^+) & 1 SO_4^{2-} to balance charges












d. magnesium hydroxide

Need 1 Mg^{2+} & 2 (OH^-) to balance charges

e. tin (IV) sulfite

Need 1 Sn^{4+} (IV means 4+ charge) & 2 (SO_3^{2-}) to balance charges

f. aluminium oxide

Need 2 Al^{3+} & 3 O^{2-} to balance charges**Oxvanions:** - a series of polyatomic ions that contains different number of oxygen atoms.**Example 2:** Name the following oxyanions.a. BrO_4^-  BrO_3^-  BrO_2^-  BrO^- b. IO_4^-  IO_3^-  IO_2^-  IO^- **Example 3:** Name the following ionic compounds.a. Na_2S Na^+ = sodium
 S^{2-} = sulfideb. KMnO_4 K^+ = potassium
 MnO_4^- = permanganatec. $\text{Li}_2\text{S}_2\text{O}_3$ Li^+ = lithium
 $\text{S}_2\text{O}_3^{2-}$ = thiosulfated. CuCr_2O_7 Cu^{2+} = copper (II)
 $\text{Cr}_2\text{O}_7^{2-}$ = dichromatee. $\text{Fe}_3(\text{BO}_3)_2$ Fe^{2+} = iron (II)
 BO_3^{3-} = boratef. $\text{Ti}(\text{BrO}_3)_3$ Ti^{3+} = titanium (III)
 BrO_3^- = bromate

Hydrate: - ionic compounds sometimes come with water molecule locked in their crystal form.
 - naming contains the ionic compound name with a prefix follow by the word “hydrate”.

Prefixes for Hydrates

| | | | |
|----------|-----------|-----------|-----------|
| 1 - mono | 4 - tetra | 7 - hepta | 10 - deca |
| 2 - di | 5 - penta | 8 - octa | |
| 3 - tri | 6 - hexa | 9 - nona | |

Example: $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ copper (II) sulfate pentahydrate

Assignment

2.8A pg. 77 #59 to 64

2.8B: Naming Simple Molecular Compounds and Acids

Nomenclature of Molecular Compounds

1. Do NOT use charges to balance subscripts. Use prefixes to name or write the formula's subscripts.
2. If the first element has one atom in the molecule, do NOT use *mono~* as a prefix.
3. The last element uses the suffix *-ide*.

Prefixes for Binary Molecular Compounds

| | | | |
|----------|-----------|-----------|-----------|
| 1 - mono | 4 - tetra | 7 - hepta | 10 - deca |
| 2 - di | 5 - penta | 8 - octa | |
| 3 - tri | 6 - hexa | 9 - nona | |

Example 1: Name the following molecular compounds.

- a. CO 1 Carbon and 1 Oxygen
Carbon monoxide
- b. CO_2 1 Carbon and 2 Oxygen
Carbon dioxide
- c. N_2O_4 2 Nitrogen and 4 Oxygen
dinitrogen tetraoxide

Example 2: Provide the chemical formula for the following compounds

- a. sulfur trioxide
 1 S and 3 O \Rightarrow **SO_3**
- b. diphosphorus pentaoxide
 2 P and 5 O \Rightarrow **P_2O_5**
- c. silicon dioxide
 1 Si and 2 O \Rightarrow **SiO_2**

Common Names for Some Molecular Compounds (Memorize!)

| | | | | | | | |
|-------------------------------------|---------|---|-------------------|------------------------|----------|---------------------------------|---------|
| H_2O | Water | H_2O_2 | Hydrogen Peroxide | O_3 | Ozone | CH_4 | Methane |
| C_3H_8 | Propane | NH_3 | Ammonia | CH_3OH | Methanol | $\text{C}_2\text{H}_5\text{OH}$ | Ethanol |
| $\text{C}_6\text{H}_{12}\text{O}_6$ | Glucose | $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ | Sucrose | | | | |

Note: Do NOT use prefixes for the above common molecular compounds!

2.8A: Naming Simple Ionic Compounds**Nomenclature:** - a naming system**IUPAC:** - International Union of Pure and Applied Chemistry.


- an organisation that oversees the standard nomenclature of all chemicals.

Nomenclature of Ionic Compounds


1. Balance the **Cation** and **Anion** Charges.
2. Use **brackets** for **multiple Complex Ions (Polyatomic Ions)**.
3. When naming, use **-ide** for the **non-metal anions**.
4. **Metals** that can have **two or more different charges** must use **Roman Numerals** in the names.

Example 1: Write the chemical formulas for the following.


a. sodium chloride

Need 1 Na^+ & 1 Cl^- to balance charges

b. calcium fluoride

Need 1 Ca^{2+} & 2 F^- to balance charges


c. ammonium sulfate

2 (NH_4^+) & 1 SO_4^{2-} to balance charges








d. magnesium hydroxide

Need 1 Mg^{2+} & 2 (OH^-) to balance charges

e. tin (IV) sulfite

Need 1 Sn^{4+} (IV means 4+ charge) & 2 (SO_3^{2-}) to balance charges

f. aluminium oxide

Need 2 Al^{3+} & 3 O^{2-} to balance charges**Oxyanions:** - a series of polyatomic ions that contains different number of oxygen atoms.**Example 2:** Name the following oxyanions.**Example 3:** Name the following ionic compounds.a. Na_2S Na^+ = sodium
 S^{2-} = sulfide**sodium sulfide**b. KMnO_4 K^+ = potassium
 MnO_4^- = permanganate**potassium permanganate**c. $\text{Li}_2\text{S}_2\text{O}_3$ Li^+ = lithium $\text{S}_2\text{O}_3^{2-}$ = thiosulfate**lithium thiosulfate**d. CuCr_2O_7 Cu^{2+} = copper (II) $\text{Cr}_2\text{O}_7^{2-}$ = dichromate**copper (II) dichromate**e. $\text{Fe}_3(\text{BO}_3)_2$ Fe^{2+} = iron (II) BO_3^{3-} = borate**iron (II) borate**f. $\text{Ti}(\text{BrO}_3)_3$ Ti^{3+} = titanium (III) BrO_3^- = bromate**titanium (III) bromate**

Chapter 3: Stoichiometry**3.1: Atomic Masses**

Average Atomic Mass Unit: - Average Mass of an atom and its isotopes taken account of their proportion of abundance (as stated on the Periodic Table of Elements).

Relative Abundance: - the relative proportion of various isotopes of an element.

Mass Spectrometer: - an instrument that measures the relative abundance of an element.
 - use magnetic and electric fields to deflect different charged isotopes. The heavier the isotope, the bigger the deflection radius.
 - the result can be graphed into a mass spectrum

$$F_{\text{centripetal}} = F_{\text{magnetic}}$$


$$\frac{mv^2}{r} = qvB$$

$$m = \frac{qvBr}{v^2}$$

$$m = \frac{qBr}{v}$$

$$m \propto r$$

q = Charge in Columb (C)
 v = speed of charged particle (m/s)
 B = magnetic field in Tesla (T)
 r = radius of deflection (m)



Example 1: State the Average Atomic Mass Unit for hydrogen if it is made of 99.48% of ^1_1H , 0.24% of ^2_1H , and 0.28% of ^3_1H .

$$\text{Average amu of Hydrogen} = \underbrace{(0.9948)(1)}_{99.48\% \text{ of Atomic Mass 1}} + \underbrace{(0.0024)(2)}_{0.24\% \text{ of Atomic Mass 2}} + \underbrace{(0.0028)(3)}_{0.28\% \text{ of Atomic Mass 3}}$$

$$\text{Average amu of Hydrogen} = 1.008$$

Example 2: Lithium has two naturally occurring isotopes. Lithium-6 has an atomic mass of 6.015 amu; lithium-7 has an atomic mass of 7.016 amu. The average atomic mass of lithium is 6.941 amu. Determine whether lithium-6 or lithium-7 is the naturally occurring atom and its percentage of abundance.

Let x = the proportion of lithium-6

Then $(1 - x)$ = proportion of lithium-7

Note: Since the average atomic mass of lithium is 6.941, which is closer to the amu of lithium-7, we can assume that the most abundant isotope is lithium-7.

$$\text{Average amu of Lithium} = (x)(6.015) + (1 - x)(7.016)$$

$$6.941 = 6.015x + 7.016 - 7.016x \quad (\text{Multiply into binomial})$$

$$6.941 - 7.016 = 6.015x - 7.016x \quad (\text{Combine Like-Terms})$$

$$-0.075 = -1.001x \quad (\text{Solve for } x)$$

$$\frac{-0.075}{-1.001} = x$$

$$x = 0.07492507 \quad (\text{Need to round to 4 significant digits})$$

Therefore, lithium-6 occurs 7.493%.

Whereas, lithium-7 occurs $(100\% - 7.493\%) = 92.51\%$

3.2: The Mole

Mole (mol): - a group of atoms or molecules numbered 6.022×10^{23} (Avogadro's Number)

Example: 1 mol of carbon (C) = 6.022×10^{23} carbon atoms = 12.01 g (same as the amu)

1 mol of oxygen (O₂) = 6.022×10^{23} oxygen molecules = 32.00 g (include subscripts with amu)

Example 1: Calculate the mass of 250 atoms of gold.

$$1 \text{ mol Au} = 196.97 \text{ g Au} = 6.022 \times 10^{23} \text{ Au atoms}$$

$$250 \text{ atoms} \times \frac{196.97 \text{ g}}{6.022 \times 10^{23} \text{ atoms}}$$

$$8.18 \times 10^{-20} \text{ g of Au}$$

Example 2: Determine the number of molecules for 50.0 mg of oxygen.

$$1 \text{ mol O}_2 = 32.00 \text{ g O}_2 = 6.022 \times 10^{23} \text{ O}_2 \text{ molecules}$$

$$(16.00 \times 2)$$

$$50.0 \text{ mg} = 0.0500 \text{ g}$$

$$0.0500 \text{ g} \times \frac{6.022 \times 10^{23} \text{ molecules}}{32.00 \text{ g}} = 9.41 \times 10^{20} \text{ molecules of O}_2$$

3.3: Molar Mass

Molar Mass (g/mol): - the mass per one mole of atoms or molecules.

- molar mass of a mono-atomic element is the same as the atomic mass.
- molar mass of a compound, binary element, or polyatomic element is the same as the combine atomic masses of all atoms in the molecule.

Example 1: Find the molar mass of the following.

a. aluminium

$$\text{Al} = 26.98 \text{ g/mol}$$

b. nitrogen

$$\text{N}_2 = 2 \times 14.01 \text{ g/mol}$$

$$\text{N}_2 = 28.02 \text{ g/mol}$$

c. water

$$\text{H}_2\text{O} = (2 \times 1.01) + 16.00$$

$$\text{H}_2\text{O} = 18.02 \text{ g/mol}$$

d. lead (IV) nitrate

$$\text{Pb}(\text{NO}_3)_4 = 207.21 + (4 \times 14.01) + (12 \times 16.00)$$

$$\text{Pb}(\text{NO}_3)_4 = 455.25 \text{ g/mol}$$

e. sucrose

$$\text{C}_{12}\text{H}_{22}\text{O}_{11} = (12 \times 12.01) + (22 \times 1.01) + (11 \times 16.00)$$

$$\text{C}_{12}\text{H}_{22}\text{O}_{11} = 342.34 \text{ g/mol}$$

Converting between Mass and Moles:

$$\text{Moles (mol)} = \frac{\text{Mass (g)}}{\text{Molar Mass (g/mol)}}$$

$$n = \frac{m}{M}$$

n = moles

m = mass

M = Molar mass

Example 2: Calculate the number of moles for:

a. 20.0 g of magnesium chloride

$$\text{MgCl}_2 = 24.31 + 2(35.45) \quad M = 95.21 \text{ g/mol}$$

$$n = \frac{m}{M} = \frac{20.0 \text{ g}}{95.21 \text{ g/mol}}$$

$$n = 0.210 \text{ mol}$$

b. 4.52 mg of glucose

$$\text{C}_6\text{H}_{12}\text{O}_6 = 6(12.01) + 12(1.01) + 6(16.00)$$

$$M = 180.18 \text{ g/mol}$$

$$n = \frac{m}{M} = \frac{4.52 \text{ mg}}{180.18 \text{ g/mol}}$$

$$n = 0.0251 \text{ mmol}$$

Example 3: Determine the mass of the following amount.

a. 8.52 mol of ozone

$$\text{O}_3 = 3(16.00) \quad M = 48.00 \text{ g/mol}$$

$$n = \frac{m}{M}$$

$$m = nM = (8.52 \text{ mol})(48.00 \text{ g/mol})$$

$$m = 409 \text{ g}$$

b. 24.7 mmol of phosphoric acid

$$\text{H}_3\text{PO}_4 = 3(1.01) + 30.97 + 4(16.00) \quad M = 98.00 \text{ g/mol}$$

$$n = \frac{m}{M}$$

$$m = nM = (24.7 \text{ mmol})(98.00 \text{ g/mol}) = 2420.6 \text{ mg}$$

$$m = 2.42 \times 10^3 \text{ mg} = 2.42 \text{ g}$$

Assignment

3.1 pg. 123 #21, 23, 35

3.2 pg. 123 #27, 29

3.3 pg. 123–124 #31, 33 to 51

3.4: Percent Composition of Compounds

Mass Percent: - the mass percentage of each element in a compound.

For Compound $A_xB_yC_z$ with its Molar Mass (M), the Mass Percentages are:

$$\%A = \frac{(x)(M_A)}{M} \times 100\% \quad \%B = \frac{(y)(M_B)}{M} \times 100\% \quad \%C = \frac{(z)(M_C)}{M} \times 100\%$$

Example 1: Calculate the mass percentage of sodium chromate.



Assume we have 161.98 g (1 mole) of Na_2CrO_4 , there are 2 moles of Na, 1 mole of Cr and 4 moles of O:

$$\% \text{Na} = \frac{(2 \text{ mol})(22.99 \text{ g/mol})}{161.98 \text{ g}} \times 100\% = 28.38622052 \%$$

$$\% \text{Cr} = \frac{(1 \text{ mol})(52.00 \text{ g/mol})}{161.98 \text{ g}} \times 100\% = 32.10272873 \%$$

$$\% \text{O} = \frac{(4 \text{ mol})(16.00 \text{ g/mol})}{161.98 \text{ g}} \times 100\% = 39.51105075 \%$$

$$\% \text{Na} = 28.386 \%$$

$$\% \text{Cr} = 32.103 \%$$

$$\% \text{O} = 39.511 \%$$

3.5: Determining the Formulas of a Compound


Empirical Formula: - the simplest ratio between the elements in a chemical formula.

Molecular Formula: - the actual chemical formula of a compound.

$$\text{Molecular Formula} = (\text{Empirical Formula})_n$$

where $n = \text{natural number}$

Example:



Note: Knowing the mass % of a compound allows us to find the empirical formula. To know the molecular formula, we must also know the molar mass.

Example 1: Vitamin C has a molar mass of 176.14 g/mol and contains carbon, hydrogen, and oxygen atoms. If the % mass of carbon and oxygen are 40.91% and 54.50% respectively, determine the empirical and molecular formula of vitamin C.

$$\% \text{ C} = 40.91\% \quad \% \text{ O} = 54.50\% \quad \% \text{ H} = 100\% - 40.91\% - 54.50\% = 4.59\%$$

Assume 100 g of Vitamin C. Then, there are

$$m_{\text{C}} = 100 \text{ g} \times 40.91\% = 40.91 \text{ g} \quad m_{\text{O}} = 100 \text{ g} \times 54.50\% = 54.50 \text{ g} \quad m_{\text{H}} = 100 \text{ g} \times 4.59\% = 4.59 \text{ g}$$

$$n_{\text{C}} = \frac{40.91 \text{ g}}{12.011 \text{ g/mol}} = 3.406044459 \text{ mol}_{\text{C}}$$

$$n_{\text{H}} = \frac{4.59 \text{ g}}{1.0079 \text{ g/mol}} = 4.554023217 \text{ mol}_{\text{H}}$$

$$n_{\text{O}} = \frac{54.50 \text{ g}}{16.00 \text{ g/mol}} = 3.40625 \text{ mol}_{\text{O}}$$

$$\frac{n_{\text{C}}}{n_{\text{O}}} = \frac{3.406044459 \text{ mol C}}{3.40625 \text{ mol O}} \approx \frac{1 \text{ mol C}}{1 \text{ mol O}}$$

$$\frac{n_{\text{H}}}{n_{\text{O}}} = \frac{4.554023217 \text{ mol H}}{3.40625 \text{ mol O}} \approx 1.33 = \frac{4 \text{ mol H}}{3 \text{ mol O}}$$

$$n_{\text{C}} : n_{\text{O}} = 1 : 1 \quad \longleftarrow \text{Combine Ratios} \quad \longrightarrow \quad n_{\text{H}} : n_{\text{O}} = 4 : 3$$

$$\text{Empirical Formula} = \text{C}_3\text{H}_4\text{O}_3 \quad (88.07 \text{ g/mol}) \quad \frac{\text{Actual Molar Mass}}{\text{Empirical Molar Mass}} = \frac{176.14 \text{ g/mol}}{88.07 \text{ g/mol}} = 2$$

$$\text{Molecular Formula} = \text{Empirical Formula} \times 2$$

$$\text{Molecular Formula} = \text{C}_6\text{H}_8\text{O}_6$$

OR

Another Method may be used where the Actual Molar Mass becomes the Mass of Vitamin used. Then, the Mole of each Atom is calculated to determine the Molecular Formula first.

$$n_{\text{C}} = \frac{40.91\% \times 176.14 \text{ g}}{12.011 \text{ g/mol}} \approx 6.00 \text{ mol}_{\text{C}}$$

$$n_{\text{H}} = \frac{4.59\% \times 176.14 \text{ g}}{1.0079 \text{ g/mol}} \approx 8.00 \text{ mol}_{\text{H}}$$

$$n_{\text{O}} = \frac{54.50\% \times 176.14 \text{ g}}{16.00 \text{ g/mol}} \approx 6.00 \text{ mol}_{\text{O}}$$

Molecular Formula (C₆H₈O₆) will be found first, then the **Empirical Formula (C₃H₄O₃)** will be stated.

Example 2: Octane, which contains carbon and hydrogen, is burned to produce 4.40 g of carbon dioxide gas and 2.03 g of water vapour. Determine the empirical and molecular formula of octane if it is found to have a molar mass of 114.26 g/mol from an analysis using a mass spectrometer.

$$n = \frac{4.40 \text{ g}}{44.011 \text{ g/mol}} = 0.0999750062 \text{ mol CO}_2$$

$$n_{\text{C}} = 0.0999750062 \text{ mol}$$

$$\frac{n_{\text{H}}}{n_{\text{C}}} = \frac{0.2253577415 \text{ mol H}}{0.0999750062 \text{ mol C}} \approx \frac{2.25 \text{ mol H}}{1 \text{ mol C}} = \frac{9 \text{ mol H}}{4 \text{ mol C}}$$

$$n = \frac{2.03 \text{ g}}{18.0158 \text{ g/mol}} = 0.1126788708 \text{ mol H}_2\text{O}$$

$$n_{\text{H}} = 0.1126788708 \text{ mol} \times 2 = 0.2253577415 \text{ mol}$$

$$\text{Empirical Formula} = \text{C}_4\text{H}_9 \quad (57.1151 \text{ g/mol})$$

$$\frac{\text{Actual Molar Mass}}{\text{Empirical Molar Mass}} = \frac{114.26 \text{ g/mol}}{57.1151 \text{ g/mol}} \approx 2$$

$$\text{Molecular Formula} = \text{Empirical Formula} \times 2$$

$$\text{Molecular Formula} = \text{C}_8\text{H}_{18}$$

Example 3: Cobalt (II) nitrate is a hydrate with a chemical formula of $\text{Co}(\text{NO}_3)_2 \cdot x\text{H}_2\text{O}$. When the 2.45 g of hydrate is heated, 1.54 g of residual is left behind. Determine the number of hydrate unit for cobalt (II) nitrate.

$$\text{Mass of Co}(\text{NO}_3)_2 = 1.54 \text{ g}$$

$$\text{Mass of H}_2\text{O released} = 2.45 \text{ g} - 1.54 \text{ g} = 0.91 \text{ g}$$

$$n = \frac{1.54 \text{ g}}{182.944 \text{ g/mol}} = 0.0084178765 \text{ mol Co}(\text{NO}_3)_2$$

$$n = \frac{0.91 \text{ g}}{18.0158 \text{ g/mol}} = 0.0505112179 \text{ mol H}_2\text{O}$$

$$\frac{n_{\text{H}_2\text{O}}}{n_{\text{Co}(\text{NO}_3)_2}} = \frac{0.0505112179 \text{ mol H}_2\text{O}}{0.0084178765 \text{ mol Co}(\text{NO}_3)_2} \approx \frac{6 \text{ mol H}_2\text{O}}{1 \text{ mol Co}(\text{NO}_3)_2}$$

$$\text{Molecular Formula} = \text{Co}(\text{NO}_3)_2 \cdot 6 \text{ H}_2\text{O}$$

Assignment

3.4 pg. 124 #54 to 56

3.5 pg. 123 #19, pg. 125 #61 to 65, 67 to 72

3.6: Chemical Equations

Chemical Reaction: - a process where **chemical change** has taken place.

Chemical Change: - a change where **New Substance(s)** are formed.

Five Evidences of a Chemical Change:

1. **Precipitate (New Solid) ↓** is formed.
2. **Colour Change**.
3. Presence of **Bubbles** or **New Odour** to indicate a **New Gas ↑**.
4. **Heat** is suddenly **Given off** or **Taken in**.
5. **Explosion!**

Reactants: - chemicals that goes into a reaction.

Products: - chemicals that are produced from a reaction.




Chemical Word Equation: - a chemical reaction written out in words.


States of Chemicals: - (s) solid, (l) liquid, (g) gas, (aq) aqueous – dissolved in water

There are 5 basic types of chemical reactions:


1. **Formation or Composition** (Element + Element → Compound)



2. **Deformation or Decomposition** (Compound → Element + Element)




3. **Single Replacement** (Element + Compound → Element + Compound)



4. **Double Replacement** (Compound + Compound → Compound + Compound)




5. **Hydrocarbon Combustion** (Hydrocarbon + Oxygen → Carbon Dioxide + Water)



3.7: Balance Chemical Equations

Coefficient: - the number in front of the chemical formula that indicates the number of moles, atoms or molecules involve in a chemical reaction.

- the absence of number in front of chemical means the coefficient is 1.

**To Predict Products and Balance Chemical Equations:**


1. Write the correct chemical formulas for all products and reactants with proper subscripts. The presence of metals or ionic compounds indicates that we will need to use ions and charges to form any products.
2. For hydrocarbon combustion, balance in the order of C, H, and then O. The product, H_2O , is always in gaseous form unless otherwise stated. (It's usually quite hot in a combustion.)
3. For other type of reactions, balance the equation for each type of cations and anions. Do NOT break up complex ions. Water may be written as HOH (H^+ and OH^-) in single and double replacement reactions.
4. Check with the Solubility Table (see Section 4.2) and the Table of Elements for the states of chemicals.

Example 1: Predict the product(s) along with the states, indicate the type of reaction, and balance the following chemical reactions.

- a. Sulfur trioxide gas is produced from its elements.



- b. A solid piece of zinc is immersed in an iron (III) chloride solution.



- c. Propane ($\text{C}_3\text{H}_8(\text{g})$) is burned in a gas barbecue.




- d. Chlorine gas is bubbled through a copper (II) iodide solution.




- e. Ammonia gas is decomposed into its elements.




f. Sulfuric acid is neutralized by sodium hydroxide solution.




g. Propanol ($\text{C}_3\text{H}_7\text{OH} (l)$) is accidentally ignited.



h. Lead (II) nitrate solution is reacted with chromium (III) sulfate solution.



i. Octane ($\text{C}_8\text{H}_{18} (l)$) is combusted in an automobile.



Assignment

3.7 pg. 126 #75 to 84

3.8: Stoichiometric Calculations: Amounts of Reactants and Products

Stoichiometry: - the calculation of quantities of chemicals in a chemical reaction.

Stoichiometry Quantities: - when all quantities of the chemical are consumed.

Gravimetric Stoichiometry: - stoichiometry that involves quantities of masses.

Mole Ratio: - the ratio of coefficients between the required chemical and the given chemical.


Gravimetric Stoichiometry Procedure:

1. Predict the products and balance the chemical equation.
2. Put all the information given under the appropriate chemicals and determine the molar masses of the chemical involved.

- Find the moles of the given chemical. $\left(n = \frac{m}{M}\right)$
- Find the mole of the required chemical using mole ratio.

$$\left(\text{mol of require} = \text{mol of given} \times \frac{\text{require coefficient}}{\text{given coefficient}}\right)$$
- Convert mole of the required chemical to its mass equivalence. ($m = nM$)

Example 1: Determine the mass of carbon dioxide formed when 50.0 kg of butane ($\text{C}_4\text{H}_{10(g)}$) is burned.




$$\textcircled{1} \quad n_{\text{C}_4\text{H}_{10}} = \frac{50.0 \text{ kg}}{58.123 \text{ g/mol}} = 0.8602446536 \text{ kmol C}_4\text{H}_{10}$$

$$\textcircled{2} \quad n_{\text{CO}_2} = 0.8602446536 \text{ kmol C}_4\text{H}_{10} \times \frac{8 \text{ mol CO}_2}{2 \text{ mol C}_4\text{H}_{10}} = 3.440978614 \text{ kmol CO}_2$$

$$\textcircled{3} \quad m_{\text{CO}_2} = nM = (3.440978614 \text{ kmol CO}_2)(44.011 \text{ g/mol}) \quad \quad \quad m_{\text{CO}_2} = 151 \text{ kg}$$

Example 2: Barium chloride solution was mixed with an excess sodium phosphate solution. What was the mass of barium chloride solid needed in the original solution to form 3.21 g of precipitate?



$$\textcircled{1} \quad n_{\text{Ba}_3(\text{PO}_4)_2} = \frac{3.21 \text{ g}}{601.938 \text{ g/mol}} = 0.0053327751 \text{ mol Ba}_3(\text{PO}_4)_2$$

$$\textcircled{2} \quad n_{\text{BaCl}_2} = 0.0053327751 \text{ mol Ba}_3(\text{PO}_4)_2 \times \frac{3 \text{ mol BaCl}_2}{1 \text{ mol Ba}_3(\text{PO}_4)_2} = 0.0159983254 \text{ mol BaCl}_2$$

$$\textcircled{3} \quad m_{\text{BaCl}_2} = nM = (0.0159983254 \text{ mol BaCl}_2)(208.236 \text{ g/mol}) \quad \quad \quad m_{\text{BaCl}_2} = 3.33 \text{ g}$$

3.9: Calculations Involving Limiting Reagents

Excess: - the reactant with more than enough amount for the reaction.


Limiting Reagent: - the reactant with the smaller amount (after taken account of the mole ratio) for the reaction.

Note: A limiting reagent question will always have enough information to find the moles of both reactants.

Steps to deal with Limiting Reagent Problems:

1. Assume one of the reactants is the limiting reagent and determine its mole amount.
2. Determine the mole amount of the other reactant.
3. Use the mole amount of the assumed limiting reagent and the mole ratio, calculate the mole amount of the other reactant actually needed.
4. If the mole amount of the other reactant is smaller than what is needed, then our assumption was wrong. The other reactant is the limiting reagent.
5. If the mole amount of the other reactant is bigger than what is needed, then our assumption was correct. It means that the other reactant is the excess.

Example 1: 5.00 g of phosphorus is reacted with 15.00 g of chlorine gas to produce phosphorus trichloride. Determine the mass of the product produced.



Since there is enough information to determine the moles of two reactants, we need to determine which one is the limiting reagent.

$$\textcircled{1} n_{\text{P}_4} = \frac{m}{M} = \frac{5.00 \text{ g}}{123.896 \text{ g/mol}} = 0.04035 \dots \text{ mol P}_4 \quad \textcircled{2} n_{\text{Cl}_2} = \frac{m}{M} = \frac{15.00 \text{ g}}{70.906 \text{ g/mol}} = 0.2115 \dots \text{ mol Cl}_2$$

Let's assume P_4 is the limiting reagent. Calculate the mol Cl_2 actually needed.

$$\textcircled{3} n_{\text{Cl}_2} = 0.040356428 \text{ mol P}_4 \times \frac{6 \text{ mol Cl}_2}{1 \text{ mol P}_4} = 0.2421385678 \text{ mol Cl}_2 \text{ needed}$$

But we don't have 0.2421385678 mol of Cl_2 , we only have 0.2115476828 mol of Cl_2 . Therefore, Cl_2 is the limiting reagent. (*Note: the limiting reagent is NOT always the chemical with the smaller number of moles. You have to always compare like we did above.*)

Now, we calculate the moles of PCl_3 formed by using moles of limiting reagent Cl_2 .

$$\textcircled{4} n_{\text{PCl}_3} = 0.2115476828 \text{ mol Cl}_2 \times \frac{4 \text{ mol PCl}_3}{6 \text{ mol Cl}_2} = 0.1410317885 \text{ mol PCl}_3$$

Finally, we determine the mass of PCl_3 produced.

$$\textcircled{5} m_{\text{PCl}_3} = nM = (0.1410317885 \text{ mol PCl}_3)(137.32 \text{ g/mol}) = 19.3664852 \text{ g}$$

$$m_{\text{PCl}_3} = 19.4 \text{ g}$$


Assignment

3.8 pg. 126 #85 to 90



3.9 pg. 127 #93 to 102, 109

Chapter 4: Types of Chemical Reactions and Solution Stoichiometry**4.1: Water, the Common Solvent****Solute:** - the matter that is being dissolved.**Solvent:** - the matter doing the dissolving.**Structure of Water:**


- V-Shaped:** - the two O-H bonds form 105° from each other, which leads to its polarity.
- Polar Molecule:** - unequal charged distribution due to the electron pairs around the oxygen atom.
- Strong O-H Hydrogen Bond:** - a type of hydrogen bond that is fairly strong compared to other types of intermolecular bonds (bonds between molecules).

**Reason that Water is a Common Solvent:**

- Polar Molecule:** - dissolves many ionic compounds due to its ability to attract cations and anions (electrolytes).
- *Note:* Some ionic compounds have low solubility (ability to dissolve) in water.

Hydration: - when ionic compound dissolves in water.**Example 1:** Sodium Chloride (Salt)

- Strong O-H Hydrogen Bond:** - dissolves many molecular compounds that have O-H hydrogen bonds.

Example 2: $\text{CH}_3\text{OH}_{(l)}$ (Methanol)

"Like-Dissolves-Like": - polar solvents tend to dissolve polar solutes; non-polar solvents tend to dissolve non-polar solutes

4.2: The Nature of Aqueous Solutions: Strong and Weak Electrolytes

Solubility Table: - a chart that shows the ability of various ion combinations to dissolve in water.

Solubility of Some Common Ionic Compounds in Water at 298.15 K (25°C)

| Ion | Group 1 NH ₄ ⁺ H ₃ O ⁺ (H ⁺) | ClO ₃ ⁻ NO ₃ ⁻ ClO ₄ ⁻ | CH ₃ COO ⁻ | Cl ⁻ Br ⁻ I ⁻ | SO ₄ ²⁻ | S ²⁻ | OH ⁻ | PO ₄ ³⁻ SO ₃ ²⁻ CO ₃ ²⁻ |
|--|--|--|------------------------------------|--|---|--|--|---|
| Solubility greater than or equal to 0.1 mol/L (very soluble) | all | all | most | most | most | Group 1 Group 2 NH ₄ ⁺ | Group 1 NH ₄ ⁺ Sr ²⁺ Ba ²⁺ Tl ⁺ | Group 1 NH ₄ ⁺ |
| Solubility less than 0.1 mol/L (slightly soluble) | none | none | Ag ⁺ Hg ⁺ | Ag ⁺ Pb ²⁺ Hg ⁺ Cu ⁺ Tl ⁺ | Ca ²⁺ Sr ²⁺ Ba ²⁺ Ra ²⁺ Pb ²⁺ Ag ⁺ | most | most | most |

1. **Strong Electrolytes:** - ionic compounds that **dissociate completely** into their ions and **conduct electricity very effectively**.

- Example 1:**
- All ionic compounds containing NH₄⁺, NO₃⁻, ClO₃⁻, or ClO₄⁻.
 - Strong Acids: HClO₄ (aq), HI (aq), HBr (aq), HCl (aq), H₂SO₄ (aq), and HNO₃ (aq)
 - Strong Bases: NH₄OH (aq), Sr(OH)₂ (aq), Ba(OH)₂ (aq), TlOH (aq), and Group 1 with OH⁻ (LiOH (aq), NaOH (aq), KOH (aq), RbOH (aq), CsOH (aq) and FrOH (aq))

2. **Weak Electrolytes:** - ionic compounds that **dissociate partially** into their ions and **conduct electricity poorly**.

- Example 2:**
- AgCl (s), PbCl₂ (s), HgCl (s), CuCl (s), and TlCl (s)
 - Weak Acids: HF (aq), HCH₃COO (aq), H₂SO₃ (aq), and other acids.
 - Weak Bases: Ca(OH)₂ (aq), Al(OH)₃ (aq), NH₃ (aq), and other bases.

3. **Non-Electrolytes:** - ionic compounds that **do not dissociate** (generally molecular compounds) in the solvent and **do not conduct electricity at all**.

- Example 3:** C₁₂H₂₂O₁₁ (aq) and C₆H₁₂O₆ (aq)

Assignment
4.2 pg. 180 – 181 #9, 11 to 14

4.3: The Composition of Solutions

Molarity (M): - moles of solute per volume of solution in Litres (mol/L).
- commonly referred to as **molar concentration (C)**.

| <u>Molar Concentration</u> | | |
|--|-------------------|----------------------|
| $\text{Molarity (mol/L)} = \frac{\text{moles (mol)}}{\text{Volume (L)}}$ | $C = \frac{n}{V}$ | 1 mol/L = 1 M |
| C = Molar Concentration | n = moles | V = Volume |

Example 1: 3.46 g of copper (II) nitrate is dissolved in 250.0 mL of water. Calculate the molarity of the solution formed.

$$m = 3.46 \text{ g Cu(NO}_3)_2$$

$$n = \frac{3.46 \text{ g}}{187.564 \text{ g/mol}}$$

$$n = 0.0184470367 \text{ mol Cu(NO}_3)_2$$

$$V = 250.0 \text{ mL} = 0.2500 \text{ L}$$

$$C = ?$$

$$C = \frac{n}{V} = \frac{0.0184470367 \text{ mol}}{0.2500 \text{ L}}$$

$$C = 0.0738 \text{ mol/L or } 73.8 \text{ mmol/L}$$

Example 2: Determine the mass of sodium dichromate needed for 500.0 mL of 0.0300 M.

$$V = 500.0 \text{ mL} = 0.5000 \text{ L}$$

$$C = 0.0300 \text{ mol/L}$$

$$M = 261.98 \text{ g/mol Na}_2\text{Cr}_2\text{O}_7$$

$$n = ? \quad m = ?$$

$$C = \frac{n}{V}$$

$$n = CV = (0.0300 \text{ mol/L})(0.5000 \text{ L})$$

$$n = 0.0150 \text{ mol}$$

$$n = \frac{m}{M}$$

$$m = nM = (0.0150 \text{ mol})(261.98 \text{ g/mol})$$

$$m = 3.93 \text{ g}$$

Example 3: The Palmense Mineral Water from the city of Fermo in Italy has a sodium hydrogen carbonate concentration of 5.207 mmol/L. What volume of this mineral water will contain 4.00 mg of sodium hydrogen carbonate?

$$m = 4.00 \text{ mg NaHCO}_3$$

$$n = \frac{4.00 \text{ mg}}{84.0079 \text{ g/mol}}$$

$$n = 0.0476145696 \text{ mmol NaHCO}_3$$

$$C = 5.207 \text{ mmol/L}$$

$$V = ?$$

$$C = \frac{n}{V}$$

$$V = \frac{n}{C} = \frac{0.0476145696 \text{ mmol}}{5.207 \text{ mmol/L}}$$

$$V = 0.00914 \text{ L or } 9.14 \text{ mL}$$

Dissociation: - when ionic compounds completely dissolve in water (100% soluble), the ionic bonds are severed and the ions “swim” freely in the new aqueous environment.

Chemical Dissociation Equation



$[A^{y+}]$ = Molar Concentration of Cation A^{y+}

$[B^{x-}]$ = Molar Concentration of Anion B^{x-}

Example 4: Write the chemical dissociation equation for the following ionic compounds when they dissolve in water.

a. $NaCl_{(s)}$




b. $Cu(NO_3)_2_{(s)}$




Example 5: Calculate the molar concentration for each ion when the following ionic compounds dissolve in water.

a. 0.150 M of $NaOH_{(aq)}$



b. 3.45 g of potassium phosphate in 500 mL of water.



Dilution: - the process where additional solvent is added to lower the concentration of an original solution.

Dilution

$$C_1V_1 = C_2V_2$$

C_1 = Concentration of Original Solution V_1 = Volume of Original Solution
 C_2 = Concentration of Diluted Solution V_2 = Total Volume of Diluted Solution

Example 6: Concentrated hydrochloric acid comes in 17.4 M. What is the volume of concentrated HCl_(aq) needed to obtain 250 mL of 1.50 M of HCl_(aq)?

$$C_1 = 17.4 \text{ mol/L} \quad C_1V_1 = C_2V_2 \quad V_1 = \frac{C_2V_2}{C_1} = \frac{(1.50 \text{ mol/L})(250 \text{ mL})}{(17.4 \text{ mol/L})}$$

$$V_1 = ? \quad V_1 = 21.6 \text{ mL}$$

$$C_2 = 1.50 \text{ mol/L}$$

$$V_2 = 250 \text{ mL}$$

Example 7: Determine the volume of water needed to dilute 30.0 mL of 0.500 M CuSO_{4(aq)} to 0.0750 M.

$$C_1 = 0.500 \text{ mol/L} \quad C_1V_1 = C_2V_2 \quad V_2 = \frac{C_1V_1}{C_2} = \frac{(0.500 \text{ mol/L})(30.0 \text{ mL})}{(0.0750 \text{ mol/L})}$$

$$V_1 = 30.0 \text{ mL} \quad V_2 = 200 \text{ mL}$$

$$C_2 = 0.0750 \text{ mol/L}$$

$$V_2 = ?$$

$$V_{\text{water}} = V_2 - 30.0 \text{ mL} = ? \quad V_{\text{water}} = 200 \text{ mL} - 30.0 \text{ mL} \quad V_{\text{water}} = 170 \text{ mL}$$

Assignment

4.3 pg. 181 #15 to 28

4.5: Precipitation Reaction


Precipitation Reaction: - a reaction where a precipitate (new solid) is formed as a product.

Insoluble: - when a solid does not seem to be soluble by the naked eye.

Slightly Soluble: - in reality, insoluble substances dissolve a tiny amount in water.


Examples:

a. Dissociation of a **Soluble Ionic Compound** and Ions Concentrations.




(Complete Ionic Dissociation – Strong Electrolytes)

b. Dissociation of a **Slightly Soluble Ionic Compound** and Ions Concentrations.



(Partial Ionic Dissociation – Weak Electrolytes)

c. Dissolving a **Molecular Compound** and Ions Concentrations.



(Soluble Molecular Compound – Non-electrolytes)

General Rules for Salts (Ionic Compounds) in Water (MEMORIZE!)

1. All NH_4^+ , NO_3^- , ClO_3^- , ClO_4^- salts are soluble.
2. Most CH_3COO^- , Cl^- , Br^- , and I^- salts are soluble (except Ag^+ , Hg^+ , Pb^{2+} , Cu^+ , and Tl^+).
3. Most SO_4^{2-} are soluble (except most Group IIA cations, Pb^{2+} and Ag^+).
4. Only Groups IA, IIA and NH_4^+ cations with S^{2-} salts are soluble.
5. Only Group IA, NH_4^+ , Sr^{2+} , Ba^{2+} , Tl^+ cations with OH^- salts are soluble (these are strong bases).
6. Only Group IA and NH_4^+ cations with PO_4^{3-} , SO_3^{2-} and CO_3^{2-} are soluble.

4.6: Describing Reactions in Solutions


Molecular Equation: - a chemical equation where compounds are written in their chemical formulas.

Complete Ionic Equation: - a chemical equation where all compounds that are soluble are written in the ionic components (slightly soluble compounds are not separated into ions).


Net Ionic Equation: - an ionic equation that only shows the ions responsible in forming the precipitate. **Spectator Ions** (ions that do not form the precipitate) are omitted.

Example 1: Predict all products form when an ammonium phosphate solution reacts with a calcium chloride solution. Explain the reaction in a form of a balanced


a. Molecular Equation



b. Complete Ionic Equation




c. Net Ionic Equation




Example 2: Predict all products form when sulfuric acid reacts with a lithium hydroxide solution. Explain the reaction in a form of a balanced


a. Molecular Equation



b. Complete Ionic Equation




c. Net Ionic Equation




Example 3: Predict all products form when solid aluminum reacts with a copper (II) nitrate solution. Explain the reaction in a form of a balanced

a. Molecular Equation



b. Complete Ionic Equation



c. Net Ionic Equation



(Need to write all the ions on both sides that correspond to any solid used or formed)

Assignment


4.5 pg. 182 #29, 30

4.6 pg. 182 #31 to 37

4.7: Stoichiometry of Precipitation Reactions**Steps to Solve a Precipitation Reaction:**


1. Write a balanced molecular equation. Identify the precipitate.
2. Put the given information underneath the proper chemicals. Identify the limiting reagent if any.
3. Using $n = CV$, convert all given information to moles.
4. Identify and use the information of the limiting reagent if necessary.
5. Determine the moles of precipitate form by using the mole ratio $\left(\frac{\text{Require Coefficient}}{\text{Given Coefficient}}\right)$.
6. Covert moles of precipitate to mass ($m = nM$).

Filtration: - a separation process to isolate the precipitate formed.

Filtration Set-up

Example 1: 200 mL of 0.0500 M of calcium chloride is reacted with 150 mL of 0.0600 M of ammonium phosphate.

- Determine the mass of the precipitate formed in this reaction.
- If the experimental mass of the precipitate is 1.28 g, calculate the % error. How can you interpret this result?
- Calculate the concentration of all ions in the final solution.



Since there is enough information to determine the moles of two reactants, we need to determine which one is the limiting reagent.

$$\textcircled{1} n_{\text{CaCl}_2} = CV = (0.0500 \text{ mol/L})(0.200 \text{ L}) = 0.01 \text{ mol}$$

$$\textcircled{2} n_{(\text{NH}_4)_3\text{PO}_4} = CV = (0.0600 \text{ mol/L})(0.150 \text{ L}) = 0.009 \text{ mol}$$

Let's assume $(\text{NH}_4)_3\text{PO}_4$ is the limiting reagent. Calculate the mol CaCl_2 actually needed.

$$\textcircled{3} n_{\text{CaCl}_2} = 0.01 \text{ mol } (\text{NH}_4)_3\text{PO}_4 \times \frac{3 \text{ mol CaCl}_2}{2 \text{ mol } (\text{NH}_4)_3\text{PO}_4} = 0.015 \text{ mol CaCl}_2 \text{ needed}$$

But we don't have 0.015 mol of CaCl_2 , we only have 0.01 mol of CaCl_2 . Therefore, CaCl_2 is the limiting reagent. (Note: the limiting reagent is NOT always the chemical with the smaller number of moles. You have to always compare like we did above.)

Now, we calculate the moles of $\text{Ca}_3(\text{PO}_4)_2$ formed by using moles of limiting reagent CaCl_2 .

$$\textcircled{2} n_{\text{Ca}_3(\text{PO}_4)_2} = 0.01 \text{ mol CaCl}_2 \times \frac{1 \text{ mol Ca}_3(\text{PO}_4)_2}{3 \text{ mol CaCl}_2} = 0.003333... \text{ mol Ca}_3(\text{PO}_4)_2$$


$$\textcircled{3} m_{\text{Ca}_3(\text{PO}_4)_2} = nM = (0.003333... \text{ mol Ca}_3(\text{PO}_4)_2)(310.18 \text{ g/mol})$$

$$m_{\text{Ca}_3(\text{PO}_4)_2} = 1.03 \text{ g}$$

b.

$$\% \text{ error} = \frac{|\text{Experimental} - \text{Theoretical}|}{\text{Theoretical}} \times 100\% = \frac{|1.28 \text{ g} - 1.03 \text{ g}|}{1.03 \text{ g}} \times 100\% \quad \text{\% error} = 24.3\%$$

This is a significant error. Since the experimental is much higher than the theoretical, we can say that there were a lot of impurities in the precipitate (from the excess ammonium phosphate).




4.8: Acid-Base Reactions

Acids: - a H^+ producer (Arrhenius); a proton, H^+ , donor (Brønsted-Lowry).



Bases: - a OH^- producer (Arrhenius); a proton, H^+ , acceptor (Brønsted-Lowry).

**Physical and Chemical Properties of Acid and Base**

| Acids | Bases |
|---|--|
| Taste Sour (Citric Acids). | Taste Bitter. |
| Burning Sensation (Stomach Acid). | Feels Slippery (Detergent, Degreaser). |
| Corrosive with Metals (reacts to give off $H_2(g)$). | Alkaline in Nature (NaOH, Baking Soda). |
| Red litmus remains Red; Blue litmus turns Red. | Red litmus turns Blue; Blue litmus remains Blue. |
| Bromothymol Blue turns Yellow | Bromothymol Blue turns Blue. |
| Phenolphthalein turns Colourless. | Phenolphthalein turns Pink. |
| pH < 7 | pH > 7 |



Example 5: Five unlabeled aqueous solutions were tested and the results are shown. If these substances are $C_2H_5OH_{(l)}$, $H_2SO_4_{(aq)}$, $HF_{(aq)}$, $NaNO_3_{(aq)}$, and $Ba(OH)_2_{(aq)}$, identify the unknown solutions.

| Solution | Conductivity | Red Litmus | Blue Litmus |
|----------|--------------|------------|-------------|
| 1 | High | Red | Blue |
| 2 | Low | Red | Red |
| 3 | High | Blue | Blue |
| 4 | None | Red | Blue |
| 5 | High | Red | Red |

Solution 1: Ionic (Conductivity High) and Neutral (No change in litmus) → **Neutral Ionic** → $NaNO_3_{(aq)}$

Solution 2: Slightly Ionic (Conductivity Low) and Acidic (Both litmus are Red) → **Weak Acid** → $HF_{(aq)}$


Solution 3: Ionic (Conductivity High) and Basic (Both litmus are Blue) → **Strong Base** → $Ba(OH)_2_{(aq)}$

Solution 4: Molecular (No Conductivity) and Neutral (No change in litmus) → **Molecular** → $C_2H_5OH_{(l)}$

Solution 5: Ionic (Conductivity High) and Acidic (Both litmus are Red) → **Strong Acid** → $H_2SO_4_{(aq)}$

Neutralization: - the reaction between acid and base to produce water and salt.

Example 6:




Steps to Solve a Neutralization Reaction:

1. Write a balanced molecular equation.
2. Put the given information underneath the proper chemicals.
3. Using $n = CV$, convert the given information to moles.
4. Determine the moles of the required chemical by using the mole ratio $\left(\frac{\text{Require Coefficient}}{\text{Given Coefficient}} \right)$.
5. Convert moles of the required chemical to concentration or volume $\left(C = \frac{n}{V} \text{ or } V = \frac{n}{C} \right)$.

Steps to Solve a Neutralization Reaction involving Limiting Reagent:

1. Write a balanced molecular equation.
2. Put the given information underneath the proper chemicals.
3. Convert all information to moles. Identify the limiting and excess reagent.
4. Determine the surplus number of moles of the excess reagent.
5. Use the total volume of both solutions; calculate the final concentration of the excess reagent.
6. Write the dissociation equation of the excess reagent.
7. Determine the concentration of the H^+ or OH^- ion.

Example 7: 30.0 mL of 0.0500 M of perchloric acid is mixed with 55.0 mL of 0.0200 M of barium hydroxide. Determine the final concentration of H^+ or OH^- ion present.



Let's assume Ba(OH)_2 is the limiting reagent. Calculate the mol HClO_4 actually needed.

$$\textcircled{1} n_{\text{HClO}_4} = 1.1 \text{ mmol Ba(OH)}_2 \times \frac{2 \text{ mmol HClO}_4}{1 \text{ mmol Ba(OH)}_2} = 2.2 \text{ mmol HClO}_4 \text{ needed}$$


But we don't have 2.2 mmol of HClO_4 , we only have 1.5 mmol of HClO_4 . Therefore, HClO_4 is the limiting reagent. (Note: the limiting reagent is NOT always the chemical with the smaller number of moles. You have to always compare like we did above.)

$$[\text{OH}^-] = \frac{1.1 \text{ mmol} - \frac{1}{2}(1.5 \text{ mmol})}{(30 \text{ mL} + 55 \text{ mL})} \quad \textcircled{[\text{OH}^-] = 0.0412 \text{ mol/L}} \quad \textcircled{[\text{H}^+] = 0 \text{ mol/L}}$$

(H^+ is a Limiting Reagent – all used up to form H_2O)

Titration: - a volumetric analysis that involves measuring the volume of known concentration solution to measure a given volume of an unknown concentration solution.

Titration Set-up



Titrant: - the solution of known concentration.

Buret: - a precise apparatus to deliver the titrant.
- the volume of the titrant added is read by subtracting the final volume and the initial volume.

Buret Valve: - can be adjusted to let one drop out at a time.

Erlenmeyer Flask: - a container commonly uses to hold the analyte. (Narrow mouth prevents splash and spillage.)

Analyte: - the solution of an unknown concentration.
- the exact volume is usually delivered by a pipet.

Acid-Base Titration: - volumetric analysis that assist in determining the unknown concentration in an acid and base neutralization.

Equivalent Point (Stoichiometric Point): - a point where the number of moles of H^+ is equivalent to the number of moles of OH^- . ($n_{H^+} = n_{OH^-}$)

Endpoint: - a point where the indicator actually changes colour to indicate neutralization is completed.

Indicator: - a chemical that changes colour due to the pH of the solution.

Common Acid-Base Indicators:

- Bromothymol Blue – Green at pH = 7
- Phenol Red – Light Orange at pH = 7
- Phenolphthalein – Light Pink at pH = 9

Example 7: Use the following observation table to determine the concentration of sulfuric acid.

| 10.0 mL of H ₂ SO ₄ (aq) titrated by 0.0350 mol/L of KOH (aq) | | | | |
|---|----------|----------|----------|----------|
| | Trial 1 | Trial 2 | Trial 3 | Trial 4 |
| Initial Volume | 0.32 mL | 24.19 mL | 3.48 mL | 24.97 mL |
| Final Volume | 24.19 mL | 45.71 mL | 24.97 mL | 46.47 mL |
| Volume of KOH added | 23.87 mL | 21.52 mL | 21.49 mL | 21.50 mL |
| Bromothymol Blue Colour | Blue | Green | Green | Green |

First, we have to complete the table by subtracting the final and the initial volumes. Since the titration is completed when the indicator turns green, we only average the result of the last 3 trials.

$$\text{Average Volume of KOH added} = \frac{21.52 \text{ mL} + 21.49 \text{ mL} + 21.50 \text{ mL}}{3} = 21.50 \text{ mL}$$



$$\textcircled{1} n_{\text{KOH}} = CV = (0.0350 \text{ mol/L}) (21.50 \text{ mL}) = 0.7525 \text{ mmol}$$

$$\textcircled{2} n_{\text{H}_2\text{SO}_4} = 0.7525 \text{ mmol KOH} \times \frac{1 \text{ mol H}_2\text{SO}_4}{2 \text{ mol KOH}} = 0.37625 \text{ mmol H}_2\text{SO}_4$$

$$\textcircled{3} C_{\text{H}_2\text{SO}_4} = \frac{n}{V} = \frac{0.37625 \text{ mmol}}{10.0 \text{ mL}} = 0.037625 \text{ mol/L}$$

$$[\text{H}_2\text{SO}_4] = 0.0376 \text{ mol/L}$$

Assignment

4.7 pg. 182–183 #39 to 44

4.8 pg. 183 #45 to 56